

# State Space Models on Special Manifolds

筑瀬靖子 (香川大学名誉教授)

We consider the state space models assuming the matrix Langevin noise processes on the Stiefel and Grassmann manifolds. The estimation of states via posterior modes is suggested.

The Stiefel manifold  $V_{k,m}$  is the space a point of which is a set of  $k$  orthonormal vectors in  $R^m$  ( $k \leq m$ ), so that  $V_{k,m} = \{X(m \times k); X'X = I_k\}$ , where  $I_k$  is the  $k \times k$  identity matrix. For  $m = k$ ,  $V_{k,m}$  is the orthogonal group  $O(m)$  of  $m \times m$  orthonormal matrices. A random matrix  $X$  on  $V_{k,m}$  is said to have the matrix Langevin (or von Mises-Fisher) distribution, denoted by  $L(m, k; F)$ , if its density function is given by (Downs [2])

$$\exp[\text{tr}(F'X)] / {}_0F_1\left(\frac{1}{2}m; \frac{1}{4}F'F\right), \text{ with } F \text{ an } m \times k \text{ matrix,}$$

where the  ${}_pF_q$  is a hypergeometric function with matrix argument. Here, assuming the rank of  $F$  being  $k$  for the simplicity of argument, we write the singular value decomposition of  $F$  as

$$F = \Gamma \Lambda \Theta', \text{ with } \Gamma \in V_{k,m}, \Theta \in O(k), \text{ and } \Lambda = \text{diag}(\lambda_1, \dots, \lambda_k), \quad \lambda_1 \geq \dots \geq \lambda_k > 0.$$

which is also expressed as  $\Gamma \Theta' \cdot \Theta \Lambda \Theta' = M \cdot C$ , say. The distribution has the unique modal orientation  $M = \Gamma \Theta' \in V_{k,m}$  and the  $\lambda_i$ 's control the concentrations about the mode in the directions determined by the orientations  $\Gamma$  and  $\Theta$ . These parameters, mode and concentrations, of the matrix Langevin distribution may be considered as the counterparts of the parameters, mean and variance-covariances, of the (multivariate) normal distribution.

The Grassmann manifold  $G_{k,m-k}$  is the space whose points are  $k$ -planes  $\nu$ , that is,  $k$ -dimensional hyperplanes in  $R^m$  containing the origin. To each  $k$ -plane  $\nu$  in  $G_{k,m-k}$ , corresponds a unique  $m \times m$  orthogonal projection matrix  $P$  idempotent of rank  $k$  onto  $\nu$ . Let  $P_{k,m-k}$  denote the set of all  $m \times m$  orthogonal projection matrices idempotent of rank  $k$ . We shall conduct our statistical analysis on the manifold  $P_{k,m-k}$  which is equivalent to the Grassmann manifold  $G_{k,m-k}$ .

For the special case  $k = 1$ , the observations from the unit hypersphere  $V_{1,m}$  are directed unit vectors, i.e., directions, and those from the real projective space  $G_{1,m-1}$

are axes or undirected lines through the origin, i.e., one-dimensional subspaces. There exists a large literature of applications of these directional statistics and its statistical analysis. The analysis of data on the general Stiefel manifold  $V_{k,m}$  is required in particular for  $k \leq m \leq 3$  in practical applications in the Earth Sciences, Medical Sciences, Astronomy, Biology, Meteorology, and other fields. See Downs [2], Watson [5], Fisher, Lewis and Embleton [3], and Mardia and Jupp [4]. One is naturally interested in  $k$ -dimensional subspaces as observations from the general Grassmann manifold  $G_{k,m-k}$ . The Grassmann manifold is a rather new subject treated as a statistical sample space. See Chikuse [1] for statistical analyses on the Stiefel and Grassmann manifolds.

We develop a state space model relating the time series observations  $\{Y_1, Y_2, \dots, Y_t\}$  on the Stiefel manifold  $V_{k,m}$  to a sequence of unobserved state modal orientation matrices  $\{X_0, X_1, \dots, X_t\}$  on  $V_{k,m}$  of noise processes distributed as matrix Langevin. We show a Bayes method for estimating the states  $\{X_1, X_2, \dots, X_t\}$  by the posterior modes assuming  $X_0$  given. An iterative procedure for the estimation is suggested. Further, we consider an extended state space model on Stiefel manifolds, where two sequences of unobserved state modal orientation matrices  $\{X_0, X_1, \dots, X_t\}$  on  $V_{k,m}$  and unobserved state regression matrices  $\{Y_0, Y_1, \dots, Y_t\}$  on  $O(m)$ , in consideration of orientational regressions, are involved.

A simple state space model on the manifold  $P_{k,m-k}$ , where the time series observations  $\{Q_1, Q_2, \dots, Q_t\}$  on  $P_{k,m-k}$  are related to a sequence of unobserved state matrices  $\{P_0, P_1, \dots, P_t\}$  on  $P_{k,m-k}$  of noise processes distributed as matrix Langevin.

A more detailed discussion is given in [Y. Chikuse, State Space Models on Special Manifolds, J. Multivariate Anal. 97 (2006) 1284–1294].

## References

- [1] Y. Chikuse, Statistics on Special Manifolds, Lecture Notes in Statistics, Vol. 174, Springer, New York, 2003.
- [2] T. D. Downs, Orientation statistics, Biometrika 59 (1972) 665–676.
- [3] N. I. Fisher, T. Lewis, B. J. J. Embleton, Statistical Analysis of Spherical Data, Cambridge University Press, London and New York, 1987.
- [4] K. V. Mardia, P. E. Jupp, Directional Statistics, Wiley, New York, 2000.
- [5] G. S. Watson, Statistics on Spheres, Lecture Notes in Mathematics (University of Arkansas) 6, Wiley, New York, 1983.